

TERMINAL 3-FOLD DIVISORIAL CONTRACTIONS OF A SURFACE TO A CURVE I

NIKOLAOS TZIOLAS

ABSTRACT. Let $\Gamma \subset X$ be a smooth curve on a 3-fold which has only index 1 terminal singularities along this curve. In this paper we investigate the existence of extremal terminal divisorial contractions $E \subset Y \longrightarrow \Gamma \subset X$, contracting an irreducible surface E to Γ . We consider cases with respect to the singularities of the general hypersurface section S of X through Γ . We completely classify the cases when S is A_i , $i \leq 3$, and D_{2n} for any n .

0. INTRODUCTION

One of the main objectives of birational geometry is to identify in each birational class of varieties some distinguished members which are “simple” and are called minimal models, and then study the structure of birational maps between them.

In dimension two, satisfactory results were known for over one hundred years. In higher dimensions, a program called the minimal model program (MMP) was developed to search for minimal models. After contributions of Reid, Mori, Kawamata, Kollár, Shokurov and others, the program was completed in dimension three by Mori in 1988. According to this program, a variety X is called a minimal model iff it is $\mathbb{Q}$ -factorial, terminal and K_X is nef. According to Mori’s theorem:

Theorem 0.1 (Mori88). *Let X be a $\mathbb{Q}$ -factorial, terminal projective 3-fold. Then there is a sequence of birational maps*

$$X = X_1 \dashrightarrow X_2 \dashrightarrow \cdots \dashrightarrow X_n = X'$$

such that X' is $\mathbb{Q}$ -factorial, terminal projective and exactly one of the following cases happen:

Received by the editors September 2001.

1991 *Mathematics Subject Classification.* Primary 14C05, 14J32; Secondary 14D15.

Key words and phrases. Algebraic geometry.

The author is supported by the Marie Curie research fellowship grant number MCFI-1999-00448.

1. $K_{X'}$ is nef and hence X' is a minimal model.
2. X' is a Mori fiber space. This means that there exists a morphism $f : X' \rightarrow S$ such that S is normal, $\dim S \leq 2$, $\rho(X'/S) = 1$ and $-K_{X'}$ is f -ample.

The birational maps that appear in the theorem are divisorial contractions and flips. It is also known [Ko-Mo98] that any birational map between minimal models is an isomorphism in codimension one and a composition of flops. Terminal flops were classified by the work of Kollár [Ko91].

The structure of birational maps between Fano fiber spaces is much more complicated. A program called the Sarkisov program was developed by Sarkisov, Reid and Corti to factorize birational maps between these spaces as a composition of so called “elementary links” [Cor95]. These links consist of flops, flips and divisorial contractions. Therefore to better understand the structure of birational maps between Fano fiber spaces, it is important to understand divisorial contractions and flips. Flips were classified by Kollár and Mori [Ko-Mo92]. However the structure of divisorial contractions is not yet completely understood.

Let $E \subset Y \xrightarrow{f} \Gamma \subset X$ be a divisorial contraction. Mori and Cutkoski completely classify such contractions when Y is Gorenstein. In particular, if $\dim \Gamma = 1$, then X is smooth along Γ and Y is just the blow up of X along Γ .

Moreover, Kawamata [Kaw94] showed that if there is a point $P \in \Gamma \subset X$ such that $P \in X$ is a cyclic quotient terminal singularity, then $\Gamma = \{P\}$ and f is a weighted blow up.

Divisorial contractions of a surface to a point, i.e. when $\Gamma = \{pt\}$, have been studied by Luo, Corti, Kawakita and others.

This paper studies divisorial contractions of a surface to a curve, i.e. when $\dim \Gamma = 1$ and X has only index 1 terminal singularities along Γ . It is not always true that given $\Gamma \subset X$, then there is a terminal contraction of a surface to Γ . We investigate when there is one, give criteria for existence or not and in the case that there is a terminal contraction we also describe the singularities of Y .

By Reid’s general elephant conjecture [Ko-Mo92], if a terminal contraction exists then there is a DuVal section $\Gamma \subset S \subset X$. We base the classification of contractions on the type of singularities of a general S through Γ instead of X itself.

Theorem 1.8 shows that under certain conditions there is always a canonical contraction.

The objective of the rest of this paper is to investigate when the contraction is terminal in the case that Γ is smooth.

I would now like to point out that for most applications this is the most interesting case and therefore it is not a big restriction to concentrate on the case that Γ is smooth. In particular, when one studies the birational rigidity of a Fano 3-fold, it is important to exclude certain curves as maximal centers. In most cases [Cor-Rei00] these curves are either lines or conics. Moreover, Miles Reid believes that a curve that is a maximal center can have at worst nodes as singularities. This is supported by a large class of examples that he calculated. Finally I believe that the methods of this paper can be used to work the case when Γ has at worst local complete intersection singularities.

In order to investigate the existence of a terminal contraction, it is important to obtain normal forms for the equations of $\Gamma \subset S \subset X$. This is done in Proposition 4.8.

Theorem 5.1 gives criteria for existence in the case that S is A_i with $i \leq 3$ and theorem 6.1 treats the case when S has D_{2n} singularities.

There is an important difference between the D_{2n} and D_{2n+1} , as well as for the higher A_n cases. The main difficulty is the explicit calculation of the $\mathbb{Q}$ -factorialization Z of E_1 , as appears in the proof of theorem 1.8. The reason of this difficulty becomes clear in lemma 6.2. I have developed a method of calculating Z and under one technical restriction i was succesfull in obtaining results for the D_{2n+1} cases too. Unfortunately at this moment I do not see how to remove this restriction or how to present the results in a “nice” and not too technical way. I plan to address these problems in a follow up paper.

Part of this work was done during my visit to Princeton University in April 2001.

I would like to thank János Kollár for many interesting and useful discussions during my visit to Princeton, and Miles Reid who suggested this problem to me.

1. UNIQUENESS AND CANONICAL CONTRACTIONS

First we define divisorial contractions.

Definition 1.1. *A 3-fold divisorial contraction is a morphism*

$$f : E \subset Y \longrightarrow \Gamma \subset X$$

such that:

1. X and Y are $\mathbb{Q}$ -factorial.
2. $Y - E \cong X - \Gamma$ and E is a prime divisor.
3. $-K_Y$ is f -ample.
4. $\rho(Y/X) = 1$.

The next proposition shows that under certain conditions, the contraction if it exists is unique.

Proposition 1.2. *Let $f : E \subset Y \longrightarrow \Gamma \subset X$ be a 3-fold divisorial contraction of an irreducible surface E to a curve Γ . Suppose that X, Y are normal, $\dim f(Y^{sing}) = 0$, X has isolated singularities and $-E$ is f -ample. Then*

$$Y \cong Proj \oplus_{d \geq 0} \mathcal{I}_{\Gamma, X}^{(d)}$$

Proof. Since $-E$ is f -ample, it follows that

$$Y \cong \oplus_{d \geq 0} f_* \mathcal{O}_Y(-dE).$$

The proposition will follow once it is shown that $f_* \mathcal{O}_Y(-dE) = \mathcal{I}_{\Gamma, X}^{(d)}$. Since by assumption X has isolated singularities and $\dim f(Y^{sing}) = 0$, there are finitely many points $P_1, \dots, P_k$ in X so that $X - \{P_1, \dots, P_k\}$ and $Y - f^{-1}\{P_1, \dots, P_k\}$ are smooth. Hence by Mori's classification of smooth terminal contractions, f is just the blow up of Γ away from $P_1, \dots, P_k$. Hence

$$f_* \mathcal{O}_Y(-dE) |_{X - \{P_1, \dots, P_k\}} = \mathcal{I}_{\Gamma, X}^d |_{X - \{P_1, \dots, P_k\}}$$

Now let $s \in \mathcal{O}_X$. Since $\dim f^{-1}(x) \leq 1, \forall x \in X$,

$$s \in f_* \mathcal{O}_Y(-dE) \iff s |_{X - \{P_1, \dots, P_k\}} \in \mathcal{I}_{\Gamma, X}^d |_{X - \{P_1, \dots, P_k\}}.$$

The proposition will now follow immediately from the following simple property of symbolic powers.

Lemma 1.3. *Let A be a ring and $\mathfrak{m}_1, \dots, \mathfrak{m}_k$ be maximal ideals. Let $I \subset A$ be a prime ideal such that $I \neq \cap_{j=1}^k \mathfrak{m}_j$. Let $a \in A$ such that*

$$a |_{Spec A - \{\mathfrak{m}_1, \dots, \mathfrak{m}_k\}} \in I^d.$$

Then $a \in I^{(d)}$.

Proof. $Spec A - \{\mathfrak{m}_1, \dots, \mathfrak{m}_k\} = \cup_{s \in \cap \mathfrak{m}_i} Spec A_s$. The assumption implies that $a \in I^d A_s, \forall s \in \cap_i \mathfrak{m}_i$. Let $s \in \cap_i \mathfrak{m}_i - I$. Then $s^\nu a \in I^d$, for some ν . Hence $a \in I^{(d)}$. $\square$

$\square$

In particular, terminal contractions of a surface to a curve are unique. However this is not true for canonical if the condition $\dim f(Y^{sing}) = 0$ is removed, as shown by the next example:

Example 1.4. *Let X be given by $x^2 + y^2 z + z^3 + t^5 = 0$ and $\Gamma : x = z = t = 0$. Then there are 2 non-isomorphic canonical divisorial contractions $g_i : S_i \subset Z_i \longrightarrow \Gamma \subset X$ contracting the surfaces $S_i \subset Z_i$*

to Γ . Z_1 has index 1 and is singular along a section of $S_1 \rightarrow \Gamma$ and Z_2 has index 2 and its singular locus is $g_2^{-1}(0)$.

Proof. The statement about the contraction $Z_2 \rightarrow X$ follows from Theorem 6.1. So here we will only show how to construct the contraction $g_1 : Z_1 \rightarrow X$.

Let $I = (x^2, z, t)$. Let $g_1 : Z_1 = B_I X \rightarrow X$ be the blow up of the ideal I in X . $Z_1 \subset \mathbb{C}^4 \times \mathbb{P}^2$. Let u, v, w be coordinates for $\mathbb{P}^2$. Look at the chart $w \neq 0$. Z_1 is given by

$$\begin{aligned} x^2 - ut &= 0 \\ u + y^2v + v^3t^2 + t^4 &= 0 \end{aligned}$$

There is only one g_1 -exceptional divisor S_1 given by $x = t = u + y^2u = 0$. Z_1 is easily seen to be singular along the line $l : x = t = u = v = 0$ which lies over Γ . Moreover, since Z_1 is a complete intersection, it has index 1. $\square$

The singularities of the general hyperplane section S of X through Γ are very important for the study of terminal contractions as shown by the following theorem.

Theorem 1.5 ([Ko-Mo92]). *Let $f : X \supset C \rightarrow Y \ni Q$ be an extremal neighborhood. Then the general member E_X of $|-K_X|$ and $E_Y = f(E_X) \in |-K_Y|$ have only DuVal singularities. Moreover, the minimal resolution of E_Y dominates E_X .*

From this it follows that if a terminal contraction exists, then there is a DuVal section S containing Γ . The converse is not true as shown by the next example.

Example 1.6. *Let X be given by $x^2 + y^2z + z^n + t^m = 0$, with $n, m \geq 4$, and $\Gamma : x = z = t = 0$. Then the contraction exists but it is only canonical, even though the section S given by $t = 0$ is D_{n+1} .*

Proof. The section S through Γ given by $t = 0$ is D_{n+1} , $\Gamma \subset S$ is of type DF_l (look at definition 4.3) and therefore by Lemma 7.3 so is the general section. Moreover, by lemma 7.4, there is no D_4 section of X containing Γ . Now the result follows from Theorem 6.1.2.a. $\square$

The next example shows that there are cases when there is no DuVal section containing Γ , and hence we can immediately conclude that there is no terminal contraction.

Example 1.7. *Let X be given by $x^2 + y^3 + z^3 + yt^6 = 0$ and $\Gamma : x = y = z = 0$. Then there is no DuVal section of X containing Γ and hence there is no terminal contraction. Moreover, the blow up $Y = B_\Gamma X$ of X along Γ is not even canonical.*

Proof. First observe that $0 \in X$ is a cD_4 singularity and therefore terminal. The section $t = 0$ is the surface

$$x^2 + y^3 + z^3 = 0$$

which is easily seen to have D_4 singularities.

A general hyperplane containing Γ in $\mathbb{C}^4$ is given by $x = ay + bz$. Hence the corresponding section of X , S , is

$$S : (ay + bz)^2 + y^3 + z^3 + yt^6 = 0.$$

This is not DuVal. To see this blow up the origin. In the affine chart given by $y = yt$, $z = zt$, the blow up is given by

$$(ay + bz)^2 + y^3t + z^3t + yt^5 = 0.$$

Blow up again. In the affine chart $y = yt$, $z = zt$, the blow up is given by

$$(ay + bz)^2 + y^3t^2 + z^3t^2 + yt^4 = 0.$$

This is singular along the line $ay + bz = t = 0$ and hence it is not normal. Therefore the original singularity is not DuVal. Hence there is no DuVal section of X containing Γ .

Now let $f : Y = B_\Gamma X \longrightarrow X$ be the blow up of X along Γ . In the affine chart $x = xz$, $y = yz$, Y is given by

$$Y : x^2z + y^3z^2 + z^2 + yt^6 = 0.$$

Moreover,

$$x^2z + y^3z^2 + z^2 + yt^6 = x^2z + (y^3 + 1)z^2 + yt^6 \sim x^2z + z^2 + yt^6$$

which is singular along the line $l : x = z = t = 0$. A typical section given by $y = a$ is ,

$$x^2z + z^2 + at^6 = 0.$$

Blow up the origin twice as before to get a nonnormal surface. Hence Y has a line of at best log canonical singularities.

Hence the existence of a DuVal section through Γ is important to conclude that $Y = B_\Gamma X$ is canonical in Proposition 2.7. $\square$

The previous example shows the significance of the existence of a DuVal section S of X through Γ . Therefore from now on we will assume the existence of such a section. In fact, we will study the existence of terminal contractions by considering cases with respect to the type of singularities of the general S through Γ , instead of the singularities of X itself.

To start with we will show that there is always a canonical contraction.

Theorem 1.8. *Let X be a $\mathbb{Q}$ -factorial index 1 terminal 3-fold and $\Gamma \subset X$ an irreducible curve having at worst plane curve singularities. Suppose there is a DuVal section S of X containing Γ . Then there is an extremal contraction $g : W \rightarrow X$ contracting an irreducible surface E to Γ , with W canonical and $\dim g(W^{\text{sing}}) = 0$. In particular, $R(\Gamma, X) = \bigoplus_{n \geq 0} \mathcal{I}_{\Gamma, X}^{(n)}$ is finitely generated.*

The proof of the previous theorem will be given in section 3.

2. SOME EASY LEMMAS

Definition 2.1. *Let X be a normal algebraic variety, and D a $\mathbb{Q}$ -Cartier divisor in X .*

1. *Let $P \in X$. Then the index of D at P , $\text{index}_P(D)$, is defined to be the smallest $r \in \mathbb{N} - 0$ such that rD is Cartier at P .*
2. *The global index of D in X , $\text{index}_X(D)$ is the smallest $r \in \mathbb{N} - 0$ such that rD is Cartier.*

Lemma 2.2. *Let X be a threefold. Suppose that its singular locus is an irreducible curve Γ and that it has only hypersurface singularities. Let D be a $\mathbb{Q}$ -Cartier divisor on X . Then*

$$\text{index}_P(D) = \text{index}_X(D)$$

for any point $P \in \Gamma$. Hence the index of D can be computed at any point of Γ .

Proof. The proof is almost identical to that for the case of isolated index 1 terminal singularities that appears in [Kaw88]. I am not aware of a reference for this more general case and therefore I include it for the convenience of the reader.

Let $r = \text{index}_X(D)$. Then there are finitely many possibilities for $\text{index}_P(D)$. Suppose that $r_1 \leq r_2 \leq \dots \leq r_k = r$ be these possibilities. Suppose that $r_1 = \text{index}_{P_1}(D)$. Since Γ is irreducible, $r_1 D$ is Cartier at all but finitely many points where D has index greater than r_1 . We can assume that it is only one, say P , since the result is local. Hence $\text{index}_P(D) = r$. Then $D' = r_1 D$ is Cartier everywhere except P . Let

$$\pi : W \rightarrow X$$

be the index 1 cover of D' . Hence

$$W - \pi^{-1}(P) \rightarrow X - P$$

is étale. X has hypersurface singularities and therefore by [Mil68, Th. 5.2]

$$\pi_1(X - P) = 0.$$

Hence the cover is trivial and therefore D' is Cartier. $\square$

Lemma 2.3. *Let $P \in \Gamma \subset X$. Suppose that $P \in X$ is a 3-dim index 1 terminal singularity, and that $P \in \Gamma$ is a plane curve singularity. Let $f : Y = B_\Gamma X \rightarrow X$ be the blow up of X along Γ . Then $f^{-1}(\Gamma) = E_1 + mE_2$. E_1 is an irreducible surface over Γ . If $P \in \Gamma$ is smooth, then $E_2 \cong \mathbb{P}^2$. Otherwise $E_2 \cong \mathbb{P}^2$ or $E_2 = \emptyset$.*

Proof. The result is local around P . Since $P \in X$ is an index 1 terminal singularity, P is a cDV point. Hence we can assume that $X \subset \mathbb{C}^4$. Suppose $P \in \Gamma$ is smooth. First we will show that $\dim f^{-1}(P) = 2$. Suppose not. Let $P \in S \subset X$ be a general hypersurface section transversal to Γ . Then $P \in S$ is DuVal and $S' = f_*^{-1}S$ is just the blow up of P in S . In particular, it is normal and $K_{S'} = f^*K_S$. If $\dim f^{-1}(P) = 1$, then $f^*S = S'$. In particular, S' is Cartier. Moreover, since f is generically the blow up of a smooth curve, $K_Y = f^*K_X + E_1$. Since Y is CM and S' Cartier,

$$\begin{aligned} K_{S'} &= (K_Y + S')|_{S'} = (f^*K_X + E_1 + f^*S)|_{S'} = \\ &= (f^*(K_X + S) + E_1)|_{S'} = f^*K_S + E_1|_{S'} = K_{S'} + E_1|_{S'} \end{aligned}$$

It now follows that $E_1|_{S'} = 0$, which is impossible. Hence $\dim f^{-1}(P) = 2$. Now in both cases, since Γ has at worst plane curve singularities at P , $\mathcal{I}_{\Gamma, \mathbb{C}^4}$ is generated by a regular sequence $\{g_1, g_2, g_3\}$. Hence $B_\Gamma \mathbb{C}^4 \rightarrow \mathbb{C}^4$ is the blow up of a regular sequence and thus all its fibers over Γ are isomorphic to $\mathbb{P}^2$. Hence $E_2 \cong \mathbb{P}^2$. $\square$

Lemma 2.4. *Let $P \in \Gamma \subset S$. Suppose that $P \in \Gamma$ is at worst a plain curve singularity, $\Gamma - P$ is smooth, and S is a normal and canonical surface. Let $f : S' = B_\Gamma S \rightarrow S$ be the blow up of S along Γ . Then S' is normal.*

Proof. Clearly, $S' - f^{-1}(P) \cong S - P$ since it is the blow up of a Cartier divisor. So the result is local over P . Hence we can assume that $S \subset X = \mathbb{C}^3$. Let $f : Y = B_\Gamma X \rightarrow X$ be the blow up of X along Γ , and E the f -exceptional divisor. Since $P \in \Gamma$ is a lci singularity, Y is just the blow up of a regular sequence, say $\{g_1, g_2\}$ in X . Hence

$$E \cong \mathbb{P}_\Gamma(I_{\Gamma, X}/I_{\Gamma, X}^2).$$

In particular, $\dim f^{-1}(x) \leq 1 \ \forall x \in X$. Hence since Y is the blow up of a smooth curve away from P , Y is normal. Moreover,

$$f^*S = S' + E \quad \text{and} \quad K_Y = f^*K_X + E.$$

Hence

$$K_Y + S' = f^*(K_X + S).$$

By [Ko97, Th. 7.3], the pair (X, S) is also canonical. Hence (Y, S') is also canonical and hence *plt*. By [Ko-Mo98, Th. 5.51] and [Ko97], S' is normal and canonical. $\square$

Lemma 2.5. *Let $P \in \Gamma \subset X$. Assume that Γ is smooth and that $P \in X$ is a 3-dimensional normal hypersurface singularity. Let $f : Y = B_\Gamma X \rightarrow X$ be the blow up of X along Γ . Let $P \in S \subset X$ be a general hypersurface section through P . Then*

$$f^*S = S' + (m_P X - 1)E_2$$

where $m_P S$ is the multiplicity of S at P and $f^{-1}(\Gamma) = E_1 + dE_2$ as in lemma 2.3.

Proof. Suppose that $f^*S = S' + aE_2$. $f^{-1}(\Gamma) = E_1 + dE_2$ is Cartier and $\mathcal{O}_Y(E_1 + dE_2) = \mathcal{O}_Y(-1)$. Then,

$$f^*S \cdot (E_1 + dE_2)^2 = S' \cdot (E_1 + dE_2)^2 + aE_2 \cdot (E_1 + dE_2)^2$$

Since $E_2 \cong \mathbb{P}^2$, it follows that $E_2 \cdot (E_1 + dE_2)^2 = 1$. Moreover, $S' = B_P S$ the blow up of S at P . Let F be the exceptional divisor. Then $F^2 = -m_P S = m_P X$, and $F^2 = S' \cdot (E_1 + dE_2)^2$. Moreover, $(E_1 + dE_2)^2 = E_1 \cdot (E_1 + dE_2) + dE_2 \cdot (E_1 + dE_2) = \Delta + dL$, where Δ is a section of $E_1 \rightarrow \Gamma$ and L a line in $E_2 = \mathbb{P}^2$. Hence

$$f^*S \cdot (E_1 + dE_2)^2 = S \cdot \Gamma = 1$$

and thus

$$a = m_P S - 1 = m_P X - 1$$

$\square$

Lemma 2.6. *Let $f : E \rightarrow \Gamma$ be a morphism from an irreducible surface to an irreducible curve. Suppose that $f^{-1}(P_0) = C_0$ is an irreducible curve and $P_0 \in \Gamma$ a smooth point. Let $f^{-1}(P) = \sum_i a_i C_i$ for another point. Then $C_0 \equiv \sum_i b_i C_i$ for $b_i \geq 0$.*

Proof. Let $\Gamma' \rightarrow \Gamma$ be the normalization of Γ , and $E' = \Gamma' \times_\Gamma E$. Hence there is a fiber square

$$\begin{array}{ccc} E' & \xrightarrow{p'} & E \\ g \downarrow & & \downarrow f \\ \Gamma' & \xrightarrow{p} & \Gamma \end{array}$$

Since Γ' is smooth, for any point Q over P , $g^{-1}(P) \equiv g^{-1}(Q)$. Apply p'_* to get the result. $\square$

Proposition 2.7. *Let $P \in \Gamma \subset X$. Assume that $P \in X$ is an index 1 terminal singularity and X normal. let $f : Y = B_\Gamma X \longrightarrow X$. Then:*

1. *If Γ is smooth, then Y is normal of index 1.*
2. *If there is a DuVal section $P \in \Gamma \subset S \subset X$ and Γ has at worst plane curve singularities, then Y is normal and canonical of index 1.*
3. *$K_Y = f^*K_X + E_1 + dE_2$, and $f^{-1}(\Gamma) = E_1 + dE_2$ is Cartier but E_1, E_2 are not $\mathbb{Q}$ -Cartier.*

Proof. First we will show that Y is CM and that ω_Y is invertible. The result is local around P . Since $P \in X$ is index 1 terminal, it is cDV. So we can assume that $X \subset \mathbb{C}^4 = U$. Let $W = B_\Gamma \mathbb{C}^4$. Since Γ has at worst plane curve singularities, W is the blow up of a regular sequence and therefore W is lci and hence CM and ω_W is invertible. Moreover,

$$E = f^{-1}(\Gamma) = E \cong \mathbb{P}_\Gamma(I_{\Gamma,U}/I_{\Gamma,U}^2)$$

is Cartier and irreducible. Suppose that $f^*X = Y + aE$, $a \in \mathbb{N}$. Hence $Y \subset W$ is Cartier and hence is CM and ω_Y is invertible.

Now assume that Γ is smooth. Since $P \in X$ is cDV, $m_P X = 2$. Let $P \in S \subset X$ be a general hypersurface section. Then $P \in S$ is DuVal. From lemma 2.5 it follows that

$$f^*S = S' + E_2$$

Since $E_2 \cong \mathbb{P}^2$ and $S' + E_2$ is Cartier, it follows that Y is smooth at the generic point of E_2 and hence regular in codimension 1 and therefore normal. This shows 1.

Now suppose that $f^{-1}(\Gamma) = E_1 + dE_2$. This is of course Cartier. Since $Y \subset W$ is Cartier, we can use adjunction to calculate K_Y . Since $W - f^{-1}(P)$ is the blow up of a smooth curve, it follows that

$$K_W = f^*K_U + 2E.$$

moreover, $f^*X = Y + E$. Hence

$$\begin{aligned} K_Y &= (K_W + Y) |_{Y=} = f^*(K_U + X) + E |_{Y=} = f^*K_X + E |_{Y=} \\ &= f^*K_X + E_1 + dE_2. \end{aligned}$$

If E_i were $\mathbb{Q}$ -Cartier, then for a fiber δ of f disjoint from E_2 and a line $l \subset E_2$, $\delta = al$, $a \geq 0$. Hence $0 \leq l \cdot E_1 = a\delta \cdot E_1 = a\delta \cdot (E_1 + dE_2) = -a < 0$, which is not possible. Of course one could argue that in this case the exceptional would have to be irreducible which is not the case.

Now suppose that a DuVal section $\Gamma \subset S \subset X$ exists. Then $S = X \cap H$ for some general plane in U . Therefore, $f^*H = H' + E$ and hence it follows that

$$f^*S = S' + E_1 + dE_2.$$

In particular, S' is Cartier. From lemma 2.4 it follows that S' is normal and canonical and therefore since S' is Cartier, Y is smooth at some points of E_2 and hence regular in codimension 1 and therefore normal.

Since X is terminal and S is canonical it follows that (X, S) is canonical. Adjunction for S' in Y gives that

$$K_Y + S' = f^*(K_X + S).$$

Hence the pair (Y, S') is also canonical. Since Y has index 1 and S' is DuVal, it follows that Y is also canonical. $\square$

If there is no DuVal section S containing Γ , then Y may fail to be canonical. In fact as shown by Example 1.7, it may not even be log canonical.

3. PROOF OF THEOREM 1.8

Let $f : Y = B_\Gamma X \longrightarrow X$. Then by proposition 2.7, Y is canonical and normal. Therefore by [Kaw88], there is $g : Z \longrightarrow Y$ such that g is an isomorphism in codimension 1 and $-E_1^Z = g_*^{-1}E_1$ is $\mathbb{Q}$ -Cartier and g -ample. More precisely, $R(E_1, Y) = \oplus_i \mathcal{O}_Y(-iE_1)$ is finitely generated and $Z = \text{Proj} R(E_1, Y)$. Hence $K_Z = g^*K_Y$ and Z is canonical. In fact since X is $\mathbb{Q}$ -factorial, Z is also $\mathbb{Q}$ -factorial. We want to contract $E_2^Z = g_*^{-1}E_2$ over X and obtain the required contraction. Let C_i be the g -exceptional curves. Of course $C_i \subset E_1^Z$. Then $K_Z \cdot C_i = 0$. On the other hand, for a $l \subset E_2^Z$, the birational transform of a line in $E_2 = \mathbb{P}^2$,

$$K_Z \cdot l = K_Y \cdot l = (E_1 + dE_2) \cdot l = -1 < 0.$$

Therefore, $\overline{NE}(Z/X)$ contains K_Z -negative extremal rays. Let R be one. Then $R = \mathbb{R}_+[\Delta]$ for an irreducible curve Δ . Δ must be contained in E_2^Z . Suppose not. Then it is a fiber of $E_1^Z \longrightarrow \Gamma$. Let $L = E_1 \cap E_2$. Then from lemma 2.6, $\Delta \equiv L^Z + \sum_i a_i C_i$, with $a_i > 0$. Hence in this case R cannot be extremal. If it was then $[C_i] \in R$ which is impossible. So all the K_Z -negative extremal curves are contained in E_2^Z . Now by the relative cone theorem there is a contraction $p : Z \longrightarrow W$ over X . There is a factorization:

$$\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ p \downarrow & & \downarrow f \\ W & \xrightarrow{q} & X \end{array}$$

By [Ko-Mo98, Prop. 3.38], W is also canonical and for any exceptional divisor, F , with center in $p(E_2^Z)$, $a(F, W) > 0$. Moreover by [Benv85], since Z is Gorenstein, $p : Z \rightarrow W$ is divisorial and therefore

$$q : W \rightarrow X$$

is the required contraction. $\square$

From the previous proof it becomes clear that W is terminal iff Z has only isolated terminal singularities along the g -exceptional curves C_i and away from E_2^Z . To investigate when this happens, it is important to obtain an explicit description of the $\mathbb{Q}$ -factorialization of E_1, Z . In general this is difficult. However in the cases that the general section S of X through Γ is D_{2n} or A_i with $i \leq 3$, it is possible to get such a description and therefore treat these cases completely.

4. NORMAL FORMS FOR $\Gamma \subset S \subset X$

We will study the existence of terminal contractions by considering cases with respect to the type of singularities that the general section S of X through Γ has. To do this it will be necessary to obtain normal forms for the equation of $\Gamma \subset S \subset X$.

There is a classification of pairs (S, Γ) when S is DuVal and $\Gamma \subset S$ a smooth curve.

Theorem 4.1 ([Jaf92]). *Let $0 \in \Gamma \subset S$ be a smooth curve in a DuVal surface S . Then under suitable coordinates, S is given by $g = 0$ and Γ by an ideal I as follows:*

Type	g	I	Where
A_n	$z^{n+1} - xy$	$I_k = (x - z^k, y - z^{n+1-k})$	$n \geq 1, 1 \leq k \leq \frac{n+1}{2}$
D_n	$x^2 + y^2z - z^{n-1}$	(x, z)	$n \geq 4$
D_{2n}	$x^2 + y^2z - z^{2n-1}$	$(x, y - z^{n-1})$	$n \geq 3$
D_{2n+1}	$x^2 + y^2z - z^{2n}$	$(x - z^n, y)$	$n \geq 2$
E_6	$x^2 + y^3 - z^4$	$(y, x + z^2)$	
E_7	$x^2 + y^3 + yz^3$	(x, y)	

Let $U \rightarrow S$ be the minimal resolution of S and let E_i be the exceptional curves. The different forms for the defining ideal of Γ correspond to the possible position of Γ in the fundamental cycle.

For the purpose of this paper it will be more convenient to fix the equations of Γ and vary this of S . The next lemma does this and it also shows the relation between the position of Γ in the fundamental cycle and its defining ideal. It essentially follows from [Jaf92].

Lemma 4.2. *With assumptions as before.*

1. Suppose that $0 \in S$ is A_n , and that the fundamental cycle is

$$\begin{array}{c} E_1 \\ \circ \end{array} - \begin{array}{c} E_2 \\ \circ \end{array} - \dots - \begin{array}{c} E_{n-1} \\ \circ \end{array} - \begin{array}{c} E_n \\ \circ \end{array} .$$

If Γ intersects the E_k then it is given by the ideal I_k .

2. Suppose that $0 \in S$ is D_n and that the fundamental cycle is

$$\begin{array}{c} E_{n-1} \\ \circ \\ | \\ \begin{array}{c} E_1 \\ \circ \end{array} - \begin{array}{c} E_2 \\ \circ \end{array} - \dots - \begin{array}{c} E_{n-2} \\ \circ \end{array} \\ | \\ \begin{array}{c} E_n \\ \circ \end{array} \end{array} .$$

Γ can only intersect E_1 , E_{n-1} or E_n . If it intersects E_1 then it is given by $I = (x, z)$ and S by $x^2 + y^2z + z^{n-1} = 0$. If it intersects E_{n-1} or E_n , then by an obvious change of variables:

- (a) In the D_{2n} case, S is given by $x^2 + y^2z + 2yz^n = 0$ and Γ by $I = (x, y)$.
- (b) In the D_{2n+1} case, S is given by $x^2 + y^2z + 2xz^n = 0$ and Γ by $I = (x, y)$.

Definition 4.3. Let Γ be a smooth curve on a surface S . Suppose that S has exactly one singular point on Γ which is of type D_n . Let $f : U \rightarrow S$ be its minimal resolution and $\Gamma' = f_*^{-1}\Gamma$. Then:

1. $\Gamma \subset S$ will be called of type FD_l if Γ' intersects E_1 in the minimal resolution of S .
2. $\Gamma \subset S$ will be called of type FD_r if Γ' intersects E_n or E_{n-1} .

Next we will derive the simplest possible normal forms for $0 \in \Gamma \subset S \subset X$. To do this it is necessary to obtain some properties of S .

First we will show that if $0 \in X$ is cA_n then $0 \in S$ is A_m for the general section S of X containing Γ .

Lemma 4.4. Let $0 \in \Gamma \subset X$, Γ a smooth curve and $0 \in X$ a cA_n 3-fold singularity. Then the general hyperplane section S of X containing Γ is A_m .

Proof. Since Γ is smooth, we can assume that it is given by $x = y = z = 0$, and X by

$$q(x, y, z) + t \left[\sum_k \phi_k(x, y, z) t^k \right] = 0$$

and the section given by $t = 0$ is A_n . Therefore

$$q(x, y, z) = q_2(x, y, z) + \sum_{i \geq 3} q_i(x, y, z)$$

with $q_2(x, y, z)$ a quadratic that is not a square.

A general hyperplane through Γ is $x = by + cz$. Then S is given by

$$q_2(by + cz, y, z) + \sum_{i \geq 3} q_i(by + cz, y, z) + t \left[\sum_i \phi_i(by + cz, y, z) t^i \right] = 0.$$

It's quadratic term is

$$\psi_2 = q_2(by + cz, y, z) + tl(by + cz, y, z) + at^2$$

where $l(by + cz, y, z)$ is linear. $0 \in S$ is A_n iff ψ_2 is not a square. If it is a square, then putting $t = 0$ it follows that $q_2(by + cz, y, z)$ is also a square.

Claim: If $q_2(by + cz, y, z)$ is a square for all b, c then $q_2(x, y, z)$ is also a square.

To see this suppose that

$$q_2(x, y, z) = a_1x^2 + a_2y^2 + a_3z^2 + a_4xy + a_5xz + a_6yz.$$

Then

$$\begin{aligned} q_2(by + cz, y, z) &= \\ &= (a_1b^2 + a_2 + a_4b)y^2 + (a_1c^2 + a_3 + a_5c)z^2 + (2bca_1 + a_4c + a_5b + a_6)yz. \end{aligned}$$

This is a square iff

$$4(a_1c^2 + a_3 + a_5c)(a_1b^2 + a_2 + a_4b) = (2bca_1 + a_4c + a_5b + a_6)^2 \quad \forall b, c.$$

Hence $a_1 = a_4 = a_5 = 0$ and $4a_2a_3 = a_6^2$. But then

$$q_2(x, y, z) = a_2y^2 + a_3z^2 + a_6yz$$

is a square. □

Now let $f : Y = B_\Gamma X \longrightarrow X$ as before. Let $f^{-1}(\Gamma) = E_1 + dE_2$. We want to relate d with some quantity on the general hyperplane section S of X , through Γ .

Lemma 4.5. *Let $f : Y = B_\Gamma X \longrightarrow X$. Assume that Y is canonical. Suppose that $f^{-1}(\Gamma) = E_1 + dE_2$. Then there is a DuVal section S of X containing Γ such that if $g : S' = f_*^{-1}S \longrightarrow S$, then*

$$g^{-1}(\Gamma) = \Gamma' + dE$$

and $E \cong \mathbb{P}^1$ is the g -exceptional curve. In fact this is true for the general the general section S of X through Γ .

This way we see that somehow d controls the type of the general hyperplane section through Γ . This result will be usefull later when we try to get normal forms for the equations of $0 \in \Gamma \subset S \subset X$.

Proof. Let $Q \in L = E_1 \cap E_2$ be a cDV point and let $S' \subset Y$ be general through Q . Then $Q \in S'$ is DuVal. Since S' is general it has the property that

$$S' \cap (E_1 + dE_2) = \Gamma' + dE$$

and Γ' maps to Γ . Moreover, for a general $\delta \subset E_1$,

$$\delta \cdot S' = 1$$

and hence

$$\delta \cdot (S' + E_1 + dE_2) = 0.$$

Therefore, there is a Cartier divisor S , $0 \in \Gamma \subset S \subset X$ such that

$$f^*S = S' + E_1 + dE_2.$$

By adjunction

$$K_{S'} = (K_Y + S')|_{S'} = f^*(K_X + S)|_{S'} = f^*K_S.$$

Therefore S is canonical with the required property. $\square$

The next proposition relates d with the type of singularities of S .

Proposition 4.6. *Let $P \in \Gamma \subset S$. Γ a smooth curve and $P \in S$ DuVal. Let $g : S' \rightarrow S$ be the blow up of S along Γ . Let E be the g -exceptional curve which is necessarily irreducible. Suppose that $g^{-1}(\Gamma) = \Gamma' + dE$. Let $f : U \rightarrow S$ be the minimal resolution of S , E_i the f -exceptional curves and Γ'' the birational transform of Γ in U . Then*

1. *Suppose that $P \in S$ is A_n and that Γ'' intersects E_k , $1 \leq k \leq \lfloor \frac{n+1}{2} \rfloor$. Then*
(a)

$$[f^{-1}(\Gamma)] = \Gamma'' + \sum_{i=1}^{k-1} iE_i + k \sum_{i=k}^{n-k+1} E_i + \sum_{i=1}^{k-1} (k-i)E_{n-k+i+1}.$$

- (b) $E = E_{n-k+1}$, and $d = k$.
2. *Suppose that $P \in S$ is D_n . Then Γ'' intersects one of the edges of the dual graph of S . Then*
(a) *With notation as in lemma 4.2.2, if Γ'' intersects E_1 , then $E = E_1$ and $d = 2$. Moreover S' is smooth along Γ' and*

$$[f^{-1}(\Gamma)] = \Gamma'' + 2 \sum_{i=1}^{n-2} E_i + E_{n-1} + E_n$$

- (b) *If Γ'' intersects E_{n-1} then*

- (i) If n is even, then $E = E_{n-1}$, $d = n/2$, S' is smooth along Γ' and

$$[f^{-1}(\Gamma)] = \Gamma'' + E_1 + 2 \sum_{i=2}^{n-2} E_i + \frac{n}{2} E_{n-1} + \frac{n-2}{2} E_n$$

- (ii) If n is odd, then $E = E_n$, $d = \frac{n-1}{2}$, and

$$[f^{-1}(\Gamma)] = \Gamma'' + E_1 + 2 \sum_{i=2}^{n-2} E_i + \frac{n-1}{2} E_{n-1} + \frac{n-1}{2} E_n.$$

Corollary 4.7. *Let $P \in \Gamma \subset S \subset X$, Γ smooth and $P \in X$ a cDV point. Let $f : Y = B_\Gamma X \rightarrow X$, $f^{-1}(\Gamma) = E_1 + dE_2$ and $L = E_1 \cap E_2$. If $P \in S$ is a D_{2n} singularity, then Y has finitely many singular points on L and therefore a $\mathbb{Q}$ -factorialization of E_1 is obtained by blowing up E_1 .*

Proof. From the previous proposition it follows that $S' = f_*^{-1}S$ is smooth at one point of L . Therefore since S' is Cartier, Y is also smooth at this point and hence has finitely many singularities along L . $\square$

Proof proposition 5.6. Let $Z = [f^{-1}(\Gamma)]$. This is an integral cycle. From the properties of the blow up it follows that $g^{-1}(\Gamma)$ is Cartier and hence $Z \cdot E = -1$ and $Z \cdot E_i = 0$ for all i such that $E_i \neq E$. Moreover, d is just the coefficient of E in Z .

We will only do the case when $0 \in S$ is D_n , with n odd, and Γ'' intersects E_{n-1} . The rest is similar. In fact the A_n is simpler.

Observe that since S has embedding dimension 4 and S' is the blow up of a smooth curve, it follows that $g^{-1}(0) = E = \mathbb{P}^1$. Therefore E appears with coefficient 1 in the fundamental cycle and hence it must be one of the edges.

So let $Z = \Gamma'' + \sum_i a_i E_i$, $a_i \in \mathbb{N}$.

There are three cases to be considered. Only one will give an integral cycle and this will be the answer.

Case 1: Check if $E = E_n$. The relations $Z \cdot E_n = -1$ and $Z \cdot E_i = 0$, $\forall i \neq n$ give the system of equations

$$\begin{aligned} -2a_1 + a_2 &= 0 \\ a_1 - 2a_2 + a_3 &= 0 \\ &\vdots \\ a_{n-3} - 2a_{n-2} + a_{n-1} + a_n &= 0 \\ a_{n-2} - 2a_{n-1} + 1 &= 0 \\ a_{n-2} - 2a_n &= -1 \end{aligned}$$

It is easy to see that the solution of this system is $a_i = i$ for $2 \leq i \leq n-2$, and $a_{n-1} = a_n = (n-1)/2$. This solution gives an integral cycle. Similarly we see that the cases $E = E_{n-1}$ and $E = E_1$ do not give integer cycles and hence are not possible.

Therefore,

$$[f^{-1}(\Gamma)] = \Gamma'' + E_1 + 2 \sum_{i=2}^{n-2} E_i + \frac{n-1}{2} E_{n-1} + \frac{n-1}{2} E_n,$$

$E = E_n$ and $d = (n-1)/2$. $\square$

We are now in position to get normal forms for the equations $0 \in \Gamma \subset S \subset X$ in the case that $0 \in S$ is D_n and S is general through Γ . We will not treat the general A_n case here and so I will not attempt to write normal forms in this case. However, normal forms for the case when $0 \in S$ is A_3 will be given in the proof of Theorem 5.1.

Proposition 4.8. *Suppose $0 \in \Gamma \subset S \subset X$, S general. Suppose that $0 \in S$ is D_n . Then under suitable choice of coordinates, $0 \in \Gamma \subset S \subset X$ is given by:*

1. *If $\Gamma \subset S$ is of type FD_l , then*

$$X : x^2 + y^2 z + z^{n-1} + t\phi_{\geq 2}(y, z, t) = 0, \quad I_\Gamma = (x, z, t).$$

Moreover no y^k appears in $\phi_{\geq 2}(y, z, t)$ for any k . For $n = 4$ this is the only possibility.

If $n \geq 5$ then $\phi_2 = 0$. That is X is given by

$$x^2 + y^2 z + z^{n-1} + t\phi_{\geq 3}(y, z, t) = 0$$

and again there is no y^k appears in $\phi_{\geq 3}(y, z, t)$.

2. *Suppose that $\Gamma \subset S$ is of type FD_r . Then*

- (a) *If n is even then $0 \in \Gamma \subset S \subset X$ is given by:*

$$x^2 + y^2 z + 2yz^{n/2} + t\phi_{\geq 2}(y, z, t) = 0, \quad I_\Gamma = (x, y, t).$$

Moreover, There is no z^k in $\phi_{\geq 2}(y, z, t)$ for any k .

(b) If n is odd then $0 \in \Gamma \subset S \subset X$ is given by:

(i)

$$x^2 + y^2 z + 2xz^{\frac{n-1}{2}} + t[axz^k + \phi_{\geq 2}(y, z, t)] = 0, \quad k \geq 1 \quad I_\Gamma = (x, y, t).$$

No yz or z^ν appear in $\phi_{\geq 2}(y, z, t)$ for any ν . $a \in \mathbb{Z}$.

(ii) Alternatively, the equation can take the form

$$x^2 + y^2 z + 2xz^{\frac{n-1}{2}} + t[xz\psi(z, t) + axz^{\frac{n-3}{2}} + bxt^k + \phi_{\geq 2}(y, z, t)] = 0,$$

$k \geq 1$, $I_\Gamma = (x, y, t)$ and in this case, y^2 , yz , or z^ν do not appear in $\phi_{\geq 2}(y, z, t)$, for any ν .

Sometimes it is better to have 2.b and sometimes 2.a. The existence of y^2 may complicate calculations.

Proof. We will apply the following methods:

1. The Weierstrass preparation theorem
2. The elimination of the y^{n-1} -term from the polynomial $a_n y^n + a_{n-1} y^{n-1} + \dots$ by a coordinate change $y \longrightarrow y - a_{n-1}/na_n$ when a_n is a unit.
3. Let M_1, M_2, M_3, M_4 be multiplicatively independent monomials in the variables x, y, z, t . Then any power series of the form $M_1 \cdot (\text{unit}) + M_2 \cdot (\text{unit}) + M_3 \cdot (\text{unit}) + M_4 \cdot (\text{unit})$ is equivalent to $M_1 + M_2 + M_3 + M_4$ by a suitable coordinate change $x \mapsto x \cdot (\text{unit})$, $y \mapsto y \cdot (\text{unit})$, $z \mapsto z \cdot (\text{unit})$, $t \mapsto t \cdot (\text{unit})$.

Case 1: Suppose that Γ'' intersects E_1 in the dual graph.

Then by lemma 4.2, $0 \in \Gamma \subset X$ is given by

$$x^2 + y^2 z + z^{n-1} + t\phi_{\geq 1}(x, y, z, t) = 0.$$

Apply the Weierstrass preparation theorem on x^2 to eliminate x from ϕ . Moreover, by lemma 4.4, if ϕ has linear terms, then $0 \in X$ is cA_n and therefore the general section $\Gamma \subset S \subset X$ is A_m which is not possible by our assumptions. Therefore $\Gamma \subset S \subset X$ is given by

$$x^2 + y^2 z + z^{n-1} + t\phi_{\geq 2}(y, z, t) = 0,$$

and $\Gamma = (x, z, t)$.

Now suppose that

$$\phi_{\geq 2}(y, z, t) = f_{\geq 2}(y) + \Phi_{\geq 2}(y, z, t),$$

with no y^k in $\Phi_{\geq 2}(y, z, t)$. Then write $f(y) = y^k \cdot (\text{unit})$, $k \geq 2$. Then the equation of X is

$$x^2 + y^2(z + ty^{k-2} \cdot (\text{unit})) + z^{n-1} + t\Phi_{\geq 2}(y, z, t) = 0.$$

The change of variables $z \mapsto z - ty^{k-2} \cdot (\text{unit})$ will give the normal form claimed by the first part of 1.

We will get further restrictions on ϕ in the case $n \geq 5$ later.

Case 2: Suppose that Γ'' intersects E_{n-1} or E_n and that $n = 2m$.

Again by lemma 4.2, and by applying the Weierstrass preparation theorem it follows that $\Gamma \subset S \subset X$ is given by

$$x^2 + y^2z + 2yz^m + t\phi_{\geq 2}(y, z, t) = 0,$$

and $I_\Gamma = (x, y, t)$. We want to eliminate z^k as well. Suppose as before that

$$\phi_{\geq 2}(y, z, t) = f_{\geq 2}(z) + \Phi_{\geq 2}(y, z, t) = z^k \cdot (\text{unit}) + \Phi_{\geq 2}(y, z, t).$$

Claim: $k \geq m$.

If this is true, then the equation of X can be written as

$$x^2 + y^2z + 2z^m(y + tz^{k-m} \cdot (\text{unit})) + t\psi_{\geq 2}(y, z, t) = 0.$$

Then the change of coordinates $y \mapsto y - tz^{k-m} \cdot (\text{unit})$ gives the claimed normal form.

To see the claim, we will show by using lemma 4.5 that if $k < m$ then there is a D_s section $\Gamma \subset T \subset X$ with $s < n$. By that lemma, we just have to show that if $f : Y = B_\Gamma X \rightarrow X$ and $f^{-1}(\Gamma) = E_1 + dE_2$, then $d < m$.

Calculate the blow up. In the chart $x = xt$, $y = yt$, Y is given by

$$x^2t + y^2tz + 2yz^m + z^k \cdot (\text{unit}) + \Phi_{\geq 2}(yt, z, t).$$

For $t = 0$ we get that $I_{E_1} = (t, 2yz^{m-k} + (\text{unit}))$, $I_{E_2} = (t, z)$ and

$$f^{-1}(\Gamma) = E_1 + kE_2$$

and $k < m$. So we obtain 2.a.

Case 3: As in case 2 but $n = 2m + 1$.

Again by the preparation theorem the equation of $0 \in \Gamma \subset S \subset X$ takes the form

$$(1) \quad x^2 + y^2z + 2xz^m + t\phi_{\geq 2}(y, z, t) = 0$$

with $I_\Gamma = (x, y, t)$.

We want to eliminate z^k . As before, write

$$\phi_{\geq 2}(y, z, t) = f_{\geq 2}(z) + \Phi_{\geq 2}(y, z, t) = z^k \cdot (\text{unit}) + \Phi_{\geq 2}(y, z, t),$$

and for the same reasons as in the previous case it follows that $k \geq m$. Moreover, in this case, $k > m$. Because suppose that $k = m$. Then by using (1), one can calculate, just like the proof of Theorem 6.1 that if $f : Y \rightarrow X$ is the blow up of X along Γ , and $f^{-1}(\Gamma) = E_1 + mE_2$, then Y has finitely many singularities along $L = E_1 \cap E_2$. But this is not possible as follows from lemma 6.2.

Then (1) becomes

$$x^2 + y^2z + 2z^m(x + tz^{k-m} \cdot (\text{unit})) + t\Phi_{\geq 2}(y, z, t) = 0.$$

The change of variables $x \mapsto x - tz^{k-m} \cdot (\text{unit})$ and $t \mapsto t/(\text{unit})$ makes the equation of X

$$x^2 + y^2z + 2xz^m + t[axz^\nu + \phi_{\geq 2}(y, z, t)] = 0,$$

$\nu \geq 1$ and $I_\Gamma = (x, y, t)$ and no power z^k appears in $\phi_{\geq 2}(y, z, t)$.

To eliminate yz , write $\phi_{\geq 2}(y, z, t) = byz + \{\text{other}\}$ and make the change of coordinates $y \mapsto y - bt/2$. This will produce the required normal form.

Sometimes it is preferable to eliminate y^2 too. To do that write $\phi_{\geq 2}(y, z, t) = cy^2 + \{\text{other}\}$ and make the change of variables $z \mapsto z - bt$. This will produce the normal form in 2.b.ii.

The only thing left to show is the second statement of 1. for $n \geq 5$. The point is that

Claim: If $\phi_2(y, z, t) \neq 0$ then there is a D_4 section S of X through Γ .

We already know that in the DF_l case it is possible to write the equations of $\Gamma \subset S \subset X$ as

$$G = x^2 + y^2z + z^{n-1} + t\phi_{\geq 2}(y, z, t) = 0,$$

and no power of y, y^k , appears in $\phi_{\geq 2}(y, z, t)$. Let

$$\phi_{\geq 2}(y, z, t) = \phi_2(y, z, t) + \phi_{\geq 3}(y, z, t).$$

Let $\phi_2(y, z, t) = a_1z^2 + a_2t^2 + a_3yz + a_4yt + a_5zt$. Let S be the section given by $t = \lambda z$. This is given by

$$F(x, y, z) = x^2 + y^2z + z^{n-1} + \lambda z\phi_2(y, z, \lambda z) + \lambda z\phi_{\geq 3}(y, z, \lambda z) = 0,$$

and

$$\phi_2(y, z, \lambda z) = (a_1 + a_2\lambda^2 + a_5\lambda)z^2 + (a_3 + a_4\lambda)yz.$$

If $\phi_2 \neq 0$, then for general λ at least one of the coefficients of z^2 or yz is not zero.

Case 1: Both coefficients are nonzero. i.e.,

$$a = a_1 + a_2\lambda^2 + a_5\lambda \neq 0 \quad \text{and} \quad b = a_3 + a_4\lambda \neq 0.$$

Hence $\phi_2(y, z, \lambda z) = az^2 + byz$, $ab \neq 0$ and

$$F = x^2 + y^2z + z^{n-1} + \lambda z(az^2 + byz) + \lambda z\phi_{\geq 3}(y, z, \lambda z).$$

Now write

$$\phi_{\geq 3}(y, z, \lambda z) = zf_{\geq 2}(y) + z^2f_{\geq 1}(y, z, \lambda z).$$

Also write $f_{\geq 2}(y) = y^k f(y)$, $k \geq 2$. Then

$$\begin{aligned} F &= x^2 + y^2 z + y z^2 [\lambda b + \lambda y^{k-1} f(y)] + z^3 [\lambda a + z^{n-4} + \lambda f_{\geq 1}(y, z, \lambda z)] = \\ &= x^2 + y^2 z + y z^2 \cdot u_1 + z^3 \cdot u_2 \end{aligned}$$

where,

$$u_1 = \lambda b + \lambda y^{k-1} f(y) \quad \text{and} \quad u_2 = \lambda a + z^{n-4} + \lambda f_{\geq 1}(y, z, \lambda z)$$

are units. Hence

$$F = x^2 + z[y^2 + yz \cdot u_1 + z^2 \cdot u_2].$$

Now eliminate the yz -term by the change of variables $y \mapsto y - zu_1/2$. Hence the equation becomes

$$F = x^2 + z[y^2 - \frac{z^2 u_1^2}{4} + z^2 u_2] = x^2 + y^2 z + z^3(u_2 - \frac{u_1^2}{4}).$$

We will now consider cases with respect to the nature of $\delta = u_2 - \frac{u_1^2}{4}$. Suppose that δ is a unit. Then

$$F \sim x^2 + y^2 z + z^3$$

and therefore S is D_4 .

Suppose now that δ is not a unit. First check when this happens. By looking at how u_1, u_2 are defined, we see that

$$\delta = (\lambda a - \frac{\lambda^2 b^2}{4}) + \{\text{higher}\}.$$

Therefore δ is not a unit iff

$$a - \frac{\lambda b^2}{4} = 0, \quad \forall \lambda.$$

But this implies that

$$a_1 + \lambda^2 a_2 + \lambda a_5 = \frac{1}{4} \lambda (a_3 + a_4 \lambda)^2$$

for all λ , and hence

$$a_1 = a_2 = a_4 = 4a_5 - a_3^2 = 0.$$

Therefore

$$\phi_2(y, z, t) = \frac{1}{4} a^2 z t + a y z.$$

Then

$$\begin{aligned}
G &= x^2 + y^2z + z^{n-1} + t\left(\frac{1}{4}a^2zt + ayz\right) + t\phi_{\geq 3}(y, z, t) = \\
&= x^2 + z\left(y^2 + \frac{1}{4}a^2t^2 + ayt\right) + z^{n-1} + t\phi_{\geq 3}(y, z, t) = \\
&= x^2 + z\left(y + \frac{1}{2}at\right)^2 + z^{n-1} + t\phi_{\geq 3}(y, z, t).
\end{aligned}$$

Make the change of variables $y \mapsto y - \frac{1}{2}at$ to bring the equation in the form

$$x^2 + y^2z + z^{n-1} + t\phi_{\geq 3}(y, z, t) = 0,$$

which is what we want.

We must now check what happens if $a = 0$ or $b = 0$ for all λ .

Suppose that $a = 0$. Hence $a_1 = a_2 = a_5 = 0$ and therefore

$$\phi_2(y, z, t) = a_3yz + a_4yt.$$

Then

$$\phi_2(y, z, \lambda z) = (a_3 + a_4\lambda)yz$$

which is nonzero for general λ . Let S again be the section defined by $t = \lambda z$. This is given by

$$\begin{aligned}
F &= x^2 + y^2z + z^{n-1} + \lambda z\phi_2(y, z, \lambda z) + \lambda z\phi_{\geq 3}(y, z, \lambda z) = \\
&= x^2 + y^2z + z^{n-1} + \lambda z^2y(a_3 + a_4\lambda) + \lambda z\phi_{\geq 3}(y, z, \lambda z) = \\
&= x^2 + z(y^2 + \lambda(a_3 + a_4\lambda)yz) + z^{n-1} + \lambda z\phi_{\geq 3}(y, z, \lambda z) = 0.
\end{aligned}$$

Now eliminate the yz term by the change of variables

$$y \mapsto y - \lambda(a_3 + a_4\lambda)z/2$$

to get

$$F = x^2 + z(y^2 - \gamma^2 z^2) + z^{n-1} + \lambda z\phi_{\geq 3}(y, z, \lambda z) = 0$$

where $\gamma = \lambda(a_3 + a_4\lambda)z/2$. Since there is no isolated power y^k it is possible to write

$$\phi_{\geq 3}(y, z, \lambda z) = zf_{\geq 2}(y) + z^2f_{\geq 1}(y, z)$$

and $f_{\geq 2}(y) = y^k f(y)$ for $k \geq 2$. Hence

$$\begin{aligned}
F &= x^2 + y^2z - \gamma^2 z^3 + z^{n-1} + \lambda z^2 y^k f(y) + \lambda z^3 f_{\geq 1}(y, z) = \\
&= x^2 + y^2z[1 + \lambda z y^{k-2} f(y)] + z^3[-\gamma + \{\text{higher}\}] = \\
&= x^2 + y^2z \cdot (\text{unit}) + z^3 \cdot (\text{unit}) \sim x^2 + y^2z + z^3,
\end{aligned}$$

and hence S is D_4 which is not possible.

The last case to consider is when $a_3 = a_4 = 0$. In this case,

$$\phi_2(y, z, t) = a_1 z^2 + a_2 t^2 + a_5 zt,$$

and therefore

$$\phi_2(y, z, \lambda z) = \gamma z^2$$

where $\gamma = a_1 + a_2 \lambda^2 + a_5 \lambda$. Hence

$$F = x^2 + y^2 z + z^{n-1} + \lambda z(\gamma z^2 + \phi_{\geq 3}(y, z, \lambda z)).$$

Again write $\phi_{\geq 3} = zy^k f(y) + z^2 f_{\geq 1}(y, z)$, and hence

$$\begin{aligned} F &= x^2 + y^2 z + z^{n-1} + \lambda z[\gamma z^2 + zy^k f(y) + z^2 f_{\geq 1}(y, z)] = \\ &= x^2 + y^2 z + z^{n-1} + \gamma \lambda z^3 + \lambda z^2 y^k f(y) + \lambda z^3 f_{\geq 1}(y, z) = \\ &= x^2 + y^2 z[1 + \lambda z f(y)] + z^3[\gamma \lambda + \lambda z^{n-3} + \lambda f_{\geq 1}(y, z)] = \\ &= x^2 + y^2 z \cdot (\text{unit}) + z^3 \cdot (\text{unit}) \sim x^2 + y^2 z + z^3, \end{aligned}$$

and therefore S is again D_4 which is impossible.

This concludes the proof of the proposition. $\square$

5. THE A_1 , A_2 , AND A_3 CASES.

In this section we will study the existence of terminal contractions in the case when $0 \in S$ is A_1 , A_2 , A_3 and Γ'' intersects the edge of the dual graph.

Theorem 5.1. *Let $P \in \Gamma \subset S \subset X$. Suppose that $0 \in S$ is A_n . Then*

1. *Suppose that Γ'' intersects one edge of the dual graph. (In particular, this is always the case when $n = 1, 2$). Then there is always a terminal contraction $E \subset W \longrightarrow \Gamma \subset X$ contracting a surface E to Γ . Let $i(W)$ be the index of W . Then*

$$n + 1 \leq i(W) \leq 2n.$$

In particular if $n = 1$ then $i(W) = 2$ and the index 2 points of W are cA . If $n = 2$ then $i(W) \in \{3, 4\}$. Moreover,

$$R(\Gamma, X) = \oplus_{d \geq 0} \mathcal{I}_{\Gamma, X}^{(d)}$$

is finitely generated by elements of degrees at most $2n$.

2. *Suppose that $n = 3$. Then*
 - (a) *If Γ'' intersects one edge of the dual graph, then there is a terminal contraction as follows from the previous part.*
 - (b) *If Γ'' intersects E_2 , i.e. the middle of the dual graph, then write the equation of $\Gamma \subset S \subset X$ as*

$$x^2 + y^2 + f_{\leq 3}(y, z, t) + f_{\geq 4}(y, z, t) = 0,$$

no y^2 appears in $f_{\leq 3}(y, z, t)$ and $I_\Gamma = (x, y, t)$. (This is always possible). Then a terminal contraction exists iff there is no proper irreducible component of $(f_{\leq 3}(y, z, t) = 0) \subset \mathbb{C}^3$ that goes through the origin. If it exists, then it has index 2 and its index 2 points are cA .

Proof. Fix notation as in the proof of Theorem 1.8.

Case 1: Assume that Γ'' intersects one edge of the dual graph.

Now proceed as in the proof of Theorem 1.8. To determine whether or not the contraction is terminal, it suffices to check whether Z has isolated terminal singularities away from E_2^Z , or not. From proposition 4.6 it follows that $d = 1$ and therefore

$$f^{-1}(\Gamma) = E_1 + E_2.$$

E_1, E_2 are both smooth and hence $Y = B_\Gamma X$ is cA_* along $L = E_1 \cap E_2$. Let $C \subset Z$ be a g -exceptional curve. It must lie over a cDV point and therefore Z can have at most finitely many terminal singularities along C . Hence W is terminal.

Now to find the index of W . Let $b \in \mathbb{N}$ such that

$$K_Z = p^*K_W + bE_2^Z.$$

Claim: E_2^Z has index n .

Let $S' = f^*S$. Then S' has exactly 1 singular point which is A_{n-1} . This follows since Γ'' intersects the edge of the dual graph. So at the generic point of L , Y has a A_{n-1} point as follows from lemma 4.5 and proposition 4.6. At this point, E_1, E_2 correspond to two lines at the edge of the dual graph. Therefore nE_1, nE_2 are Cartier at all but finitely many points.

Since $E_1 + E_2$ is Cartier, it follows that the singularities of Y lie on $L = E_1 \cap E_2$. In fact if $n \geq 2$, then Y is singular along L . If not then by lemma 4.5, there is a $\Gamma \subset S \subset X$ such that $S' = B_\Gamma S$ is smooth along $S' \cap E_1$. But by proposition 4.6 this implies that S must be A_1 which is not possible.

Moreover, since Z has index 1, it has hypersurface singularities and therefore by lemma 2.2 the indices of E_1^Z and E_2^Z can be computed at any point of L^Z . Therefore, they have index n .

Let $l \subset E_2^Z$ be the birational transform of a line in $E_2 = \mathbb{P}^2$ contracted by p . Then, $l \cdot E_2^Z = -a/n$, $a \in \mathbb{N}$. Moreover,

$$l \cdot E_2^Z + l \cdot E_1^Z = -1$$

and clearly $l \cdot E_1^Z \leq 1$. Hence

$$-2 \leq l \cdot E_2^Z \leq -1.$$

Therefore $n \leq a \leq 2n$. Moreover, $K_Z \cdot l = K_Y \cdot l = -1$ and hence $b = n/a$. In fact $a = n$ is not possible. If it was then

$$K_Z = p^* K_W + E_2^Z.$$

Moreover,

$$K_W = q^* K_X + E_1^W \quad \text{and} \quad K_Z = h^* K_X + E_1^Z + E_2^Z$$

where $h = f \circ g : Z \longrightarrow X$. Combining the above relations it follows that

$$p^* E_1^W \cong E_1^Z$$

which is not possible. Now the claim about the indices follows immediately.

The statement about the type of singularities follows from [Ko-Mo92, Theorem 4.7].

Case 2: $n = 3$. The only case to study is when Γ'' intersects the middle of the dual graph. It will be necessary to obtain a normal form for the equation $\Gamma \subset S \subset X$.

Lemma 5.2. *Let $0 \in \Gamma \subset S \subset X$. Suppose that $0 \in S$ is an A_3 singular point and that Γ'' intersects the middle of the dual graph in the minimal resolution of S . Then under suitable coordinates, $0 \in \Gamma \subset S \subset X$ can be written as*

$$x^2 + y^2 + 2xz^2 + t\phi_{\geq 1}(x, z, t) = 0$$

and $I_\Gamma = (x, y, t)$ and no power z^k appears in $\phi_{\geq 1}(x, z, t)$.

Proof. By Theorem 4.1, under suitable coordinates it is possible to write $S : xy - z^5 = 0$ and $\Gamma : x - z^2 = y - z^2 = 0$. The change of coordinates $x \mapsto x + z^2$, $y \mapsto y + z^2$ brings it to

$$xy + xz^2 + yz^2 = 0$$

and $\Gamma : x = y = 0$. Now let $x \mapsto x - y$, $y \mapsto x + y$ and apply the Weierstrass preparation theorem to y^2 to get

$$S : x^2 + y^2 + 2xz^2 + t\phi_{\geq 1}(x, z, t) = 0$$

and $\Gamma : x = y = t = 0$. To eliminate z^k we must show that z does not appear in $\phi_{\geq 1}(x, z, t)$. If it does then it is easy to see that the general hyperplane section $y = at + bx$ through Γ will be A_1 which is impossible.

So now proceed as in proposition 4.8 to eliminate powers of z . $\square$

The statement about existence of a terminal contraction is proved in exactly the same way as Theorem 6.1 and I omit its proof. $\square$

Example 5.3. Let X be given by

$$x^2 + y^2 + 2xz^2 + t^m = 0$$

and $\Gamma : x = y = t = 0$. Then there is no 3-fold terminal contraction, contracting a surface to Γ .

Proof. In this case, the section $S : (t = 0)$ is an A_3 type and the curve intersects the middle of the minimal resolution of S . Moreover, $f_{\leq 3}(x, z, t) = 2xz^2$ and there is a plane through the origin in $f_{\leq 3}(x, z, t) = 0$. Therefore by the previous theorem there is no terminal contraction. $\square$

6. THE D_{2n} CASES.

In this section we will study the existence of terminal contractions when $0 \in S$ is a D_{2n} type for general $0 \in \Gamma \subset S \subset X$.

Theorem 6.1. Let $P \in \Gamma \subset S \subset X$. Suppose that $P \in S$ is a D_n type singular point. Then

1. Suppose that $n = 4$. Write the equation of $P \in \Gamma \subset S \subset X$ as

$$x^2 + f_3(y, z, t) + f_{\geq 4}(y, z, t) = 0$$

where $I_\Gamma = (x, z, t)$ and $f_3(y, z, t)$ is homogeneous of degree 3. This is always possible. Then

- (a) If $f_3(y, z, t)$ is an irreducible homogeneous cubic, then there is a terminal contraction $W \rightarrow X$ of a surface to Γ . W has index 2 and has exactly one singularity which is of cA_x type. Moreover,

$$R(\Gamma, X) = \oplus_{d \geq 0} \mathcal{I}_{\Gamma, X}^{(d)}$$

is finitely generated by elements of degrees 1 and 2.

- (b) If $f_3(y, z, t)$ is reducible or 0, then there is no terminal contraction.
2. Suppose $n \geq 5$.
 - (a) Suppose that $\Gamma \subset S$ is of type FD_l . Then there is no terminal contraction.
 - (b) Suppose that $\Gamma \subset S$ is of type FD_r and n is even. Then write the equation of $P \in \Gamma \subset S \subset X$

$$x^2 + f_3(y, z, t) + f_{\geq 4}(y, z, t) = 0$$

with $I_\Gamma = (x, y, t)$.

- (i) If $f_3(y, z, t)$ is an irreducible homogeneous cubic, then there is a terminal contraction $W \rightarrow X$ of a surface to Γ . W

has index 2 and has exactly one singularity which is of cD type. Moreover,

$$R(\Gamma, X) = \oplus_{d \geq 0} \mathcal{I}_{\Gamma, X}^{(d)}$$

is finitely generated by elements of degrees 1 and 2.

- (ii) If $f_3(y, z, t)$ is reducible or 0, then there is no terminal contraction.

Proof. we will only do the second part of the theorem. The first one is proved in exactly the same way.

Case 1: Γ'' intersects E_1 in the dual graph and $n \geq 5$.

By proposition 4.8, under suitable coordinates $\Gamma \subset S \subset X$ is given by

$$x^2 + y^2z + z^{n-1} + t\phi_{\geq 3}(y, z, t) = 0$$

$\Gamma : x = z = t = 0$, and no power y^k appears. Let

$$f : Y = B_{\Gamma}X \longrightarrow X$$

be the blow up of X along Γ . In the chart $x = xt, z = zt$ it is given by

$$x^2t + y^2z + z^{n-1}t^{n-2} + \phi_{\geq 3}(y, zt, t) = 0.$$

For $t = 0$ we see that

$$f^{-1}(\Gamma) = E_1 + 2E_2$$

and $E_1 : z = t = 0$, $E_2 : y = t = 0$. Moreover $L = E_1 \cap E_2 : y = z = t = 0$ and it is easy to see that Y has only one singular point on L , the origin. However it is singular along the line $l : x = y = z = 0$ which lies in E_2 .

Therefore

$$g : Z = B_{E_1}Y \longrightarrow Y$$

is a $\mathbb{Q}$ -factorialization of E_1 . In the chart $z = zt$ it is given by

$$x^2 + y^2z + z^{n-1}t^{2n-4} + \frac{1}{t}\phi_{\geq 3}(y, zt^2, t) = 0.$$

Write

$$\phi_{\geq 3}(y, zt^2, t) = tf_{\geq 2}(y) + t^2f_{\geq 1}(y, z, t).$$

It is easy to see that $f_{\geq 1}(0, z, 0) = 0$. Then Y is given by

$$x^2 + y^2z + z^{n-1}t^{2n-4} + f_{\geq 2}(y) + tf_{\geq 1}(y, z, t) = 0.$$

Now let $C = g^{-1}(0) : x = y = t = 0$. Then, since $f_{\geq 1}(0, z, 0) = 0$ it is clear that Z is singular along C . Moreover since $E_2^Z \cong E_2 \cong \mathbb{P}^2$, it is clear that C is not contained in E_2^Z .

Now contract $E_2^Z \cong \mathbb{P}^2$ over X to get a contraction

$$p : W \longrightarrow X.$$

However W is only canonical and not terminal since it is singular along C . Therefore in this case there is no terminal contraction.

Case 2: Γ'' intersects the other edge of the dual graph and $n = 2m$ is even.

Then again by proposition 4.8, under suitable coordinates $\Gamma \subset S \subset X$ is given by

$$x^2 + y^2z + 2yz^m + t\phi_{\geq 2}(y, z, t) = 0,$$

$\Gamma : x = y = t = 0$ and no power z^k appears. The method is the same as before. Blow up Γ . In the chart $x = xt, y = yt, Y$ is given by

$$x^2t + y^2zt + 2yz^m + \phi_{\geq 2}(yt, z, t) = 0.$$

For $t = 0$ we find that

$$f^{-1}(\Gamma) = E_1 + mE_2$$

and $E_1 : y = t = 0, E_2 : z = t = 0$. As before, Y has exactly one singular point on $L = E_1 \cap E_2 = (y, z, t)$. Let

$$\phi_{\geq 2}(y, z, t) = \phi_2(y, z, t) + \phi_{\geq 3}(y, z, t).$$

Whether or not W is terminal depends on what kind of singularities Z has away from E_2^Z . In the chart $y = yt, Z$ is given by

$$x^2 + y^2t^2z + 2yz^m + \frac{1}{t}\phi_2(yt^2, z, t) + \frac{1}{t}\phi_{\geq 3}(yt^2, z, t) = 0.$$

Write again

$$\phi_{\geq 3}(yt^2, z, t) = tf_{\geq 2}(z) + t^2f_{\geq 1}(y, z, t)$$

with $f_{\geq 1}(y, 0, 0) = 0$. Then Z is given by

$$F = x^2 + y^2t^2z + 2yz^m + \frac{1}{t}\phi_2(yt^2, z, t) + f_{\geq 2}(z) + tf_{\geq 1}(y, z, t) = 0.$$

Let $C = g^{-1}(0) : x = z = t = 0$, and

$$\Phi_2(y, z, t) = \frac{1}{t}\phi_2(yt^2, z, t).$$

We want to investigate the singularities of Z along C .

$$\begin{aligned}\frac{\partial F}{\partial x} &= 2x \\ \frac{\partial F}{\partial y} &= 2yzt^2 + 2z^m + \frac{\partial \Phi_2}{\partial y} + t \frac{\partial f_{\geq 1}}{\partial y} \\ \frac{\partial F}{\partial z} &= y^2t^2 + 2myz^{m-1} + \frac{\partial \Phi_2}{\partial z} + \frac{\partial f_{\geq 2}}{\partial z} + t \frac{\partial f_{\geq 1}}{\partial z} \\ \frac{\partial F}{\partial t} &= 2y^2tz + \frac{\partial \Phi_2}{\partial t} + f_{\geq 1}(y, z, t) + t \frac{\partial f_{\geq 1}}{\partial t}\end{aligned}$$

Hence Z is singular along a point $Q \in C$ iff

$$\frac{\partial \Phi_2}{\partial y}(Q) = \frac{\partial \Phi_2}{\partial z}(Q) = \frac{\partial \Phi_2}{\partial t}(Q) = 0.$$

Let

$$\phi_2(y, z, t) = a_1y^2 + a_2t^2 + a_3yt + a_4yz + a_5zt.$$

Then

$$\Phi_2(y, z, t) = \frac{1}{t}\phi_2(yt^2, z, t) = a_1y^2t^3 + a_2t + a_3yt^2 + a_4yzt + a_5z.$$

Now it follows that

$$\begin{aligned}\frac{\partial \Phi_2}{\partial y} &= 2a_1yt^3 + a_3t^2 + a_4zt \\ \frac{\partial \Phi_2}{\partial z} &= a_4yt + a_5 \\ \frac{\partial \Phi_2}{\partial t} &= 3a_1y^2t^2 + a_2 + 2a_3yt + a_4yz\end{aligned}$$

Along C ,

$$\frac{\partial \Phi_2}{\partial y} = 0, \quad \frac{\partial \Phi_2}{\partial z} = a_5, \quad \frac{\partial \Phi_2}{\partial t} = a_2$$

Therefore, Z is either singular along C and $a_2 = a_5 = 0$, or has exactly one singular point in E_2^Z (in the other chart). If $a_2 = a_5 = 0$, then

$$\phi_2(y, z, t) = y(a_1y + a_3t + a_4z).$$

A coordinate independent way to say this property is the statement of 2.b.

We will now find the index of the singularities of W . Suppose that

$$K_Z = p^*K_W + aE_2^Z.$$

Moreover, $E_2^Z \cong E_2 \cong \mathbb{P}^2$. E_1^Z is Cartier and therefore for a general line $l \subset E_2^Z$, $l \cdot E_1^Z = 1$. On the other hand,

$$l \cdot (E_1^Z + 2E_2^Z) = l \cdot g^*(E_1 + 2E_2) = l \cdot (E_1 + 2E_2) = -1.$$

Hence $l \cdot E_2^Z = -1$. Moreover,

$$l \cdot K_Z = l \cdot g^* K_Y = l \cdot K_Y = -1.$$

Combining the above we see that $a = 1$ and therefore

$$K_Z = p^* K_W + E_2^Z.$$

E_2^Z has index 2 and hence W has also index 2. From the above proof it is also clear that W has exactly one index 2 point. This, as well as the type of the singularities, also follows from [Ko-Mo92, Theorem 4.7].

Moreover, since W has index 2 it follows that $-2E$ is p -very ample where $E = E_1^W$. The statement about the number of generators of $R(\Gamma, X)$ follows immediately. $\square$

The difference between the D_{2n} and D_{2m+1} cases is shown by the next lemma.

Lemma 6.2. *Suppose that the general section S of X containing Γ is D_{2n+1} and $\Gamma \subset S$ is of type DF_r . Let $f : Y = B_\Gamma X \rightarrow X$. Let $f^{-1}(\Gamma) = E_1 + dE_2$. Then Y is singular along $L = E_1 \cap E_2$.*

This is precisely the reason that makes the D_{2n+1} case very difficult to work with. In the D_{2n} cases, Y had exactly one singular point on L and that made an explicit description of the $\mathbb{Q}$ -factorialization of E_1 relatively easy.

Proof. Suppose that Y is not singular along L . Let $Q \in L$ be a smooth point. Let S' be a general section of Y through Q . As in lemma 4.4, there is a section S of X through Γ such that $S' = f_*^{-1}S$. Then by assumption, S is D_{2n+1} . Then by proposition 5.6.2.b.ii, it follows that $Q \in S'$ is singular which is not true. $\square$

The previous result could of course be proved by explicitly calculating Y by using the normal forms of Proposition 4.8.

The results of Theorem 6.1 depend on the singularities of the general section S of X containing Γ . To apply the theorem it would be useful to get information about the general section from a special section. The next lemma gives information about the general section starting from a special one.

Lemma 6.3. *Let $\Gamma \subset X$. Suppose that the general section of X containing Γ is D_k . Let $P \in \Gamma \subset S_0 \subset X$ be a special section and suppose that S_0 is D_n with $n \geq 5$. Then*

1. *If $\Gamma \subset S_0$ is of type DF_l , then the general section S of X through Γ is D_m , $m \leq n$ and also of type DF_l .*
2. *If $\Gamma \subset S_0$ is of type DF_r and n is even, then the general S through Γ is D_{2k} and also of type DF_r .*

Proof. Case 1: Suppose that $\Gamma \subset S_0$ is of type FD_l , i.e., Γ'' intersects E_1 in the fundamental cycle of S_0 . Let S be the general section through Γ and assume it is D_m . If Γ'' intersects E_{m-1} or E_m in the fundamental cycle of S , then by proposition 4.6 it follows that $d = m/2$, if m is even, or $d = (m-1)/2$, if m is odd. On the other hand, by the assumption on S_0 and Proposition 4.8, under suitable coordinates $\Gamma \subset X$ is given by

$$x^2 + y^2z + z^{n-1} + t\phi_{\geq 2}(y, z, t) = 0,$$

and $I_\Gamma = (x, z, t)$. Use notation as in lemma 6.2.

A computation as in Theorem 6.1 shows that $d = 2$ and Y has exactly one singular point on L . Hence the only possibility that the general section is not as claimed is that it is D_5 and Γ'' intersects E_4 or E_5 . But then Y is singular along L as follows from lemma 6.2.

The fact that $m \leq n$ follows from the upper semicontinuity of the Tyurina number of the singularity.

Case 2: Suppose that $\Gamma \subset S_0$ is of type FD_r and n is even. First we will show that it is not possible that $\Gamma \subset S$, $S \subset X$ general through Γ , is of type FD_l . Suppose it is. Then since n is even, $S' = f_*^{-1}S$ has exactly one singular point which is D_{n-1} as follows from proposition 4.6.2.a. On the other hand, S'_0 has exactly one A_{n-1} singular point Q . Therefore $Q \in Y$ is cA_k and by [KoBa88] it is cA_k in a neighborhood of Q . But then for a general $\Gamma \subset S \subset X$, S' is A_k and hence it must be of type FD_r .

If $\Gamma \subset S$ is FD_r but S is D_{2m+1} for general S , then by lemma 6.2 Y is singular along L which is not true as follows from corollary 4.7. $\square$

Hence by looking at one section we know in which part of theorem 6.1 we are. So if we know that there is a section as in 1 then all we need to know to conclude that there is no terminal contraction is that the general section is not D_4 . The next lemma gives a criterion for that.

Lemma 6.4. *Let $\Gamma \subset X$ be given by*

$$x^2 + f_{\geq 3}(y, z, t) = 0,$$

and $\Gamma = (x, y, t)$. Moreover suppose that $t = 0$ is a DuVal section S of X containing Γ and Γ'' intersects E_1 in the fundamental cycle of S . Then a D_4 section of X containing Γ does not exist iff

$$f_3(y, z, t) = g(y, z, t)h^2(y, z, t).$$

Proof. According to proposition 4.8, in suitable coordinates $\Gamma \subset X$ is given by

$$x^2 + y^2z + z^{n-1} + t\phi_{\geq 2}(y, z, t) = 0,$$

and $I_\Gamma = (x, z, t)$. The cubic term then of the above equation is

$$q_3(y, z, t) = y^2z + t\phi_2(y, z, t).$$

From the proof of the first part of proposition 4.8 it follows that a D_4 section exists iff

$$q_3(y, z, t) \neq g(y, z, t)h^2(y, z, t),$$

for any $g(y, z, t), h(y, z, t)$, which is the condition claimed by the lemma. $\square$

Example 6.5. *Let X be given by*

$$x^2 + y^2z + 2yz^n + t^m + t\phi_{\geq 3}(y, z, t) = 0,$$

and $I_\Gamma = (x, y, t)$. Then there is a terminal contraction iff $m = 3$.

Proof. The section S_0 given by $(t = 0)$ is D_{2n} and $\Gamma \subset S_0$ is DF_r . Hence by lemma 6.3 so is the general section through Γ , S . Now apply theorem 6.1. $\square$

Finally, I would like to mention that it will be interesting to get a more direct way of constructing the contraction of theorems 5.1, 6.1. In particular, is it true that it is a weighted blow up of the curve? I plan to address this question as well as treat the remaining cases in a future paper.

REFERENCES

- [Art66] A. Artin, *On isolated rational singularities of surfaces*, Amer. J. Math. 88 (1966), pp 129-136.
- [Benv85] X. Benveniste, *Sur le cone des 1-cycles effectifs en dimension 3*, Math. Ann. 272 (1985), pp 257-265.
- [Cor95] A. Corti, *Factoring birational maps of threefolds after Sarkisov*, J. Algebraic Geometry 4 (1995), pp 223-254.
- [Cor-Rei00] A. Corti, M. Reid, *Explicit birational geometry of 3-folds*, LMN 281 (2000)
- [Cut88a] S. Cutkosky, *Elementary contractions of Gorenstein threefolds*, Math. Ann. 280 (1988), pp 521-525.
- [Cut88b] S. Cutkosky, *Weil divisors and symbolic algebras*, Duke Math. J. 57 (1988), pp 175-183.
- [Jaf92] D. B. Jaffe, *Local geometry of smooth curves passing through rational double points*, Math. Ann. 294 (1992), pp 645-660.
- [ka01] M. Kawakita, *Divisorial contractions in dimension three which contract divisors to smooth points*, Invent. Math. 145 (2001) no 1, pp 105-119.
- [Kaw88] Y. Kawamata, *Crepant blowing up of 3-dimensional canonical singularities and its application to degenerations of surfaces*, Annals of Mathematics 127 (1988), pp 93-163.

- [Kaw94] Y. Kawamata, *Divisorial contractions to 3-dimensional terminal quotient singularities*, in Higher-dimensional complex varieties (Trento 1994), 241-246, de Gruyter, Berlin, 1996.
- [Ko97] J. Kollár, *Singularities of pairs*, Proc. Symp. Pure Math. Soc. volume 62, 1997.
- [Ko91] J. Kollár, *Flips, flops, minimal models, etc.*, Surv. in Diff. Geom. 1 (1992), pp 293-323.
- [KoBa88] J. Kollár, N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. 91 (1988), pp 299-338.
- [Ko-Mo98] J. Kollár, S. Mori, *Birational geometry of algebraic varieties*, Cambridge University Press, 1988.
- [Ko-Mo92] J. Kollár, S. Mori, *Classification of three-dimensional flips*, J. Am. Math. Soc. Vol. 5 no. 3 (1992), pp 533-703.
- [Mil68] J. Milnor, *Singular points of complex hypersurfaces*, Princeton University Press 1968.
- [Rei87] M. Reid, *Young person's guide to canonical singularities*, Proc. Symp. Pure Math. Soc. volume 46, 1987.

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL,
ENGLAND

E-mail address: `tziolas@maths.warwick.ac.uk`